

On some generalizations of Nadler's contraction principle

Coroian Iulia

Faculty of Mathematics and Computer Science, Babeş-Bolyai University Cluj-Napoca

coroian.iulia@gmail.com

The purpose of this work is to present some generalizations of the well known Nadler's contraction principle. More precisely, using an axiomatic approach of the Pompeiu-Hausdorff metric we will study the properties of the fractal operator generated by a multivalued contraction.

References

- [1] M. A. Khamsi, W.A. Kirk, *An Introduction to Metric Spaces and Fixed Point Theory, Pure and Applied Mathematics*, Wiley-Interscience, New York, 2001.
- [2] W.A. Kirk, N. Shahzad, *Remarks on metrics transform and fixed point theorems*, 2013. Article ID 9612004682934060 .
- [3] H. K. Pathak, N. Shahzad, *A new fixed point result and its application to existence theorem for nonconvex Hammerstein type integral inclusions* , Electronic Journal of Qualitative Theory of Differential Equations , No.62, 2012, 1-13.
- [4] S. B. Nadler, Jr., *Multi-valued contraction mappings*, Pacific J. Math., Vol.30, No.2, 1969, 475-487.
- [5] A. Petruşel, I. A. Rus, M. A. Marcel, *The role of eq metrics in fixed point theory*, Topol. Meth. Nonlinear Anal., 41(2013), 85-112.
- [6] I. A. Rus, *Generalized Contraction and Applications*, Cluj Univ. Press, Cluj-Napoca, 2001.
- [7] I. A. Rus, A. Petruşel, G.Petruşel, *Fixed Point Theory*, Cluj University Press, 2008.