

Hardy-Littlewood-Stein-Weiss type theorems for Riesz potentials and their commutators in Morrey spaces

Canay Aykol and Javanshir J. Hasanov

Abstract. In this paper we consider weighted Morrey spaces $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$. We prove the Hardy-Littlewood-Stein-Weiss type $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$ theorems for Riesz potential I^α and its commutators $[b, I^\alpha]$ and $|b, I^\alpha|$, where $0 < \alpha < n$, $0 \leq \lambda < n - \alpha$, $1 < p < \frac{n-\lambda}{\alpha}$, $-n + \lambda \leq \gamma < n(p-1) + \lambda$, $\mu = \frac{q\gamma}{p}$, $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$, $b \in BMO(\mathbb{R}^n)$. As a result of these we obtain the conditions for the boundedness of the commutator $|b, I^\alpha|$ from Besov-Morrey spaces $B_{p,\theta,\lambda,|\cdot|^\gamma}^s(\mathbb{R}^n)$ to $B_{q,\theta,\lambda,|\cdot|^\mu}^s(\mathbb{R}^n)$. Furthermore, we consider the Schrödinger operator $-\Delta + V$ on \mathbb{R}^n and obtain weighted Morrey $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ estimates for the operators $V^s(-\Delta + V)^{-\beta}$ and $V^s\nabla(-\Delta + V)^{-\beta}$. Finally we apply our results to various operators which are estimated from above by Riesz potentials.

Mathematics Subject Classification (2010): 42B20, 42B25, 42B35.

Keywords: Riesz potential, commutator, fractional maximal operator, Schrödinger operator, Hardy-Littlewood-Stein-Weiss type estimate, Morrey space, BMO space.

1. Introduction

The well known Morrey spaces $\mathcal{L}^{p,\lambda}(\Omega)$ introduced by Charles Morrey (see [24]) in 1938 in relation to the study of partial differential equations, and presented in various books, see e.g. [11, 16, 39]. They were widely investigated during the last decades, including the study of classical operators of harmonic analysis maximal, singular and potential operators on Morrey spaces and their various generalizations

Received 30 May 2020; Accepted 06 November 2020.

© Studia UBB MATHEMATICA. Published by Babeş-Bolyai University



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

have found wide applications in many problems of real analysis and partial differential equations. Morrey spaces are defined by the norm

$$\|f\|_{\mathcal{L}^{p,\lambda}} = \sup_{x, t>0} t^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,t))},$$

where $0 \leq \lambda < n$, $1 \leq p < \infty$ and $B(x, t)$ is the open ball in \mathbb{R}^n of radius t centered at x . In the theory of partial differential equations, together with weighted Lebesgue spaces, Morrey spaces play an important role. Later, Morrey spaces found important applications to Navier-Stokes ([22], [39]) and Schrödinger ([28], [29], [30], [33], [34]) equations, elliptic problems with discontinuous coefficients ([5], [8]), and potential theory ([1], [2]).

The results on the boundedness of potential operators and classical Calderón-Zygmund singular operators go back to [1] and [27], respectively, while the boundedness of the maximal operator in the Euclidean setting was proved in [6].

Hardy-Littlewood-Stein-Weiss inequality in the Lebesgue spaces was proved by H.G. Hardy and J.E. Littlewood [12] in the one-dimensional case and by E.M. Stein and G. Weiss [37] in the case $n > 1$. In the Lebesgue and Morrey spaces with variable exponent the Hardy-Littlewood-Stein-Weiss inequality was proved by S.G. Samko [31] and J.J. Hasanov [13], respectively.

Let f be a locally integrable function on \mathbb{R}^n . The so-called fractional maximal function is defined by the formula

$$M^\alpha f(x) = \sup_{t>0} |B(x,t)|^{-1+\alpha/n} \int_{B(x,t)} |f(y)| dy, \quad 0 \leq \alpha < n,$$

where $|B(x, t)|$ is the Lebesgue measure of the ball $B(x, t)$ such that $|B(x, t)| = \omega_n t^n$ in which ω_n denotes the volume of the unit ball in \mathbb{R}^n . It coincides with the Hardy-Littlewood maximal function $Mf \equiv M_0 f$. Maximal operators play an important role in the differentiability properties of functions, singular integrals and partial differential equations. They often provide a deeper and more simplified approach to understanding problems in these areas than other methods.

Fractional maximal operator is intimately related to the Riesz potential

$$I^\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y) dy}{|x - y|^{n-\alpha}}, \quad 0 < \alpha < n,$$

such that

$$M^\alpha f(x) \leq \omega_n^{\frac{\alpha}{n}-1} (I^\alpha |f|)(x).$$

The aim of this paper is to give the necessary and sufficient conditions for the boundedness of Riesz potential I^α and its commutators from weighted Morrey spaces $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{p,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$. We also obtain the necessary conditions for the boundedness of the commutator $|b, I^\alpha|$ from Besov-Morrey spaces $B_{p,\theta,\lambda,|\cdot|^\gamma}^s(\mathbb{R}^n)$ to $B_{q,\theta,\lambda,|\cdot|^\mu}^s(\mathbb{R}^n)$. Furthermore, we consider the Schrödinger operator $-\Delta + V$ on \mathbb{R}^n and obtain weighted Morrey $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ estimates for the operators $V^s(-\Delta + V)^{-\beta}$ and $V^s \nabla(-\Delta + V)^{-\beta}$. Finally we apply our results to various operators which are estimated from above by Riesz potentials.

Throughout the paper we use the letters c, C for positive constants, independent of appropriate parameters and not necessarily the same at each occurrence. If $A \leq CB$ and $B \leq CA$, we write $A \approx B$ and say that A and B are equivalent.

2. Preliminaries

We use the following notation. For $1 \leq p < \infty$, $L_p(\mathbb{R}^n)$ is the space of all classes of measurable functions on \mathbb{R}^n for which

$$\|f\|_{L_p} = \left(\int_{\mathbb{R}^n} |f(x)|^p dx \right)^{\frac{1}{p}} < \infty,$$

up to the equivalence of the norms

$$\|f\|_{L_p} \sim \sup_{\|g\|_{L_{p'}} \leq 1} \left| \int_{\mathbb{R}^n} f(y)g(y)dy \right| \tag{2.1}$$

and also $WL_p(\mathbb{R}^n)$, the weak L_p space defined as the set of all measurable functions f on \mathbb{R}^n such that

$$\|f\|_{WL_p} = \sup_{r>0} r |\{x \in \mathbb{R}^n : |f(x)| > r\}|^{1/p} < \infty.$$

For $p = \infty$ the space $L_\infty(\mathbb{R}^n)$ is defined by means of the usual modification

$$\|f\|_{L_\infty} = \text{ess sup}_{x \in \mathbb{R}^n} |f(x)|.$$

For $1 \leq p < \infty$ let $L_{p,\omega}(\mathbb{R}^n)$ be the space of measurable functions on \mathbb{R}^n such that

$$\|f\|_{L_{p,\omega}} = \|f\omega^{1/p}\|_{L_p(\mathbb{R}^n)} = \left(\int_{\mathbb{R}^n} |f(x)|^p \omega(x) dx \right)^{1/p} < \infty,$$

and for $p = \infty$ the space $L_{\infty,\omega}(\mathbb{R}^n) = L_\infty(\mathbb{R}^n)$.

Definition 2.1. The weight function ω belongs to the class $A_p(\mathbb{R}^n)$ for $1 \leq p < \infty$, if the following statement

$$\sup_{x \in \mathbb{R}^n, t > 0} \frac{1}{|B(x, t)|} \int_{B(x, t)} \omega(y) dy \left(\frac{1}{|B(x, r)|} \int_{B(x, r)} \omega^{-\frac{1}{p-1}}(y) dy \right)^{p-1}$$

is finite and ω belongs to $A_1(\mathbb{R}^n)$, if there exists a positive constant C such that for any $x \in \mathbb{R}^n$ and $t > 0$

$$|B(x, t)|^{-1} \int_{B(x, t)} \omega(y) dy \leq C \text{ess sup}_{y \in B(x, t)} \frac{1}{\omega(y)}.$$

The following theorem was proved in [37].

Theorem 2.2. Let $0 < \alpha < n$, $1 < p < \frac{n}{\alpha}$, $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$, $\alpha p - n < \gamma < n(p - 1)$, $\mu = \frac{\alpha\gamma}{p}$. Then the operators M^α and I^α are bounded from $L_{p,|\cdot|^{-\gamma}}(\mathbb{R}^n)$ to $L_{q,|\cdot|^{-\mu}}(\mathbb{R}^n)$.

Theorem 2.3. [36] *Let $1 < p < \infty$ and $-n < \gamma < n(p - 1)$. Then the operator M is bounded on $L_{p,|\cdot|^\gamma}(\mathbb{R}^n)$.*

Let M^\sharp be the sharp maximal function defined by

$$M^\sharp f(x) = \sup_{t>0} |B(x, t)|^{-1} \int_{B(x,t)} |f(y) - f_{B(x,t)}| dy,$$

where $f_{B(x,t)}(x) = |B(x, t)|^{-1} \int_{B(x,t)} f(y) dy$.

Definition 2.4. We define the $BMO(\mathbb{R}^n)$ space as the set of all locally integrable functions f with finite norm

$$\|f\|_{BMO} = \sup_{x \in \mathbb{R}^n, t>0} |B(x, t)|^{-1} \int_{B(x,t)} |f(y) - f_{B(x,t)}| dy$$

or

$$\|f\|_{BMO} = \inf_C \sup_{x \in \mathbb{R}^n, t>0} |B(x, t)|^{-1} \int_{B(x,t)} |f(y) - C| dy.$$

Definition 2.5. We define the $BMO_{p,\omega}(\mathbb{R}^n)$ ($1 \leq p < \infty$) space as the set of all locally integrable functions f with finite norm

$$\|f\|_{BMO_{p,\omega}} = \sup_{x \in \mathbb{R}^n, t>0} \frac{\|(f(\cdot) - f_{B(x,t)})\chi_{B(x,t)}\|_{L_{p,\omega}(\mathbb{R}^n)}}{\|\chi_{B(x,t)}\|_{L_{p,\omega}(\mathbb{R}^n)}}.$$

Theorem 2.6. [14, Theorem 4.4] *Let $1 \leq p < \infty$ and ω be a Lebesgue measurable function. If $\omega \in A_p(\mathbb{R}^n)$, then the norms $\|\cdot\|_{BMO_{p,\omega}}$ and $\|\cdot\|_{BMO}$ are mutually equivalent.*

We find it convenient to define the Morrey and weighted Morrey spaces in the form as follows.

Definition 2.7. Let $1 \leq p < \infty$. Morrey spaces $L_{p,\lambda}(\mathbb{R}^n)$ and weighted Morrey spaces $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ are defined by the norms

$$\|f\|_{L_{p,\lambda}} = \sup_{x \in \mathbb{R}^n, t>0} t^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,t))}$$

and

$$\|f\|_{L_{p,\lambda,|\cdot|^\gamma}} = \sup_{x \in \mathbb{R}^n, t>0} t^{-\frac{\lambda}{p}} \|f\|_{L_{p,|\cdot|^\gamma}(B(x,t))},$$

respectively.

For $1 \leq p, \theta \leq \infty$ and $0 < s < 1$, Besov-Morrey space $B_{p,\theta,\lambda,|\cdot|^\gamma}^s(\mathbb{R}^n)$ consists of all functions $f \in L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ such that

$$\|f\|_{B_{p,\theta,\lambda,|\cdot|^\gamma}^s} = \|f\|_{L_{p,\lambda,|\cdot|^\gamma}} + \left(\int_{\mathbb{R}^n} \frac{\|f(x - \cdot) - f(\cdot)\|_{L_{p,\lambda,|\cdot|^\gamma}}^\theta}{|x|^{n+s\theta}} dx \right)^{1/\theta} < \infty.$$

3. Riesz potential operator in the spaces $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$

In this section we prove the Hardy-Littlewood-Stein-Weiss type $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$ -theorem for Riesz potential I^α , where $-n + \lambda \leq \gamma < n(p - 1) + \lambda$, $1 < p < \frac{n-\lambda}{\alpha}$, $\mu = \frac{q\gamma}{p}$ and $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$.

First we give following theorems which we use while proving our main results.

Theorem 3.1. [25] *Let $1 < p < \infty$, then $M : L_{p,\varphi}(\mathbb{R}^n) \rightarrow L_{p,\varphi}(\mathbb{R}^n)$ if and only if $\varphi \in A_p(\mathbb{R}^n)$.*

Theorem 3.2. [15] *Let $1 < p < \infty$, $0 \leq \lambda < n$, $\varphi \in A_p(\mathbb{R}^n)$, then $M : L_{p,\lambda,\varphi}(\mathbb{R}^n) \rightarrow L_{p,\lambda,\varphi}(\mathbb{R}^n)$.*

Theorem 3.3. *Let $0 < \alpha < n$, $0 \leq \lambda < n - \alpha$, $1 < p < \frac{n-\lambda}{\alpha}$, $-n + \lambda \leq \gamma < n(p - 1) + \lambda$ and $\mu = \frac{q\gamma}{p}$. Then the operator I^α is bounded from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$ if and only if $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$.*

Proof. Sufficiency: Let $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$ and $f \in L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$. Then

$$\begin{aligned} |I^\alpha f(x)| &= \left(\int_{B(x,t)} + \int_{\mathbb{R}^n \setminus B(x,t)} \right) |f(y)| |x - y|^{\alpha-n} dy \\ &\equiv F_1(x, t) + F_2(x, t). \end{aligned}$$

First we estimate $F_1(x, t)$. By using Hölder's inequality we have

$$\begin{aligned} F_1(x, t) &= \int_{B(x,t)} |f(y)| |x - y|^{\alpha-n} dy \\ &\leq \sum_{j=-\infty}^{-1} (2^j t)^{\alpha-n} \int_{B(x,2^{j+1}t) \setminus B(x,2^j t)} |f(y)| dy \\ &\leq Ct^\alpha Mf(x). \end{aligned} \tag{3.1}$$

Now we estimate $F_2(x, t)$. By using Hölder's inequality we get

$$\begin{aligned} F_2(x, t) &\leq \int_{\mathbb{R}^n \setminus B(x,t)} |f(y)| |x - y|^{\alpha-n} dy \\ &\leq \sum_{j=0}^{\infty} (2^j t)^{\alpha-n} \int_{B(x,2^{j+1}t) \setminus B(x,2^j t)} |f(y)| dy \\ &\leq \sum_{j=0}^{\infty} (2^j t)^{\alpha-n} \|\chi_{B(x,2^{j+1}t)}\|_{L_{p'(\cdot),|\cdot|^\gamma/(1-p)}} \|f\chi_{B(x,2^{j+1}t)}\|_{L_{p,|\cdot|^\gamma}} \\ &\leq Ct^{\alpha - \frac{n-\lambda}{p}} |x|^{-\frac{\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}} \sum_{j=0}^{\infty} 2^{j(\alpha - \frac{n-\lambda}{p})} \\ &\leq Ct^{\alpha - \frac{n-\lambda}{p}} |x|^{-\frac{\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}} \end{aligned}$$

Thus

$$F_2(x, t) \leq Ct^{\alpha - \frac{n-\lambda}{p}} |x|^{-\frac{\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}. \tag{3.2}$$

Therefore from (3.1) and (3.2) we get

$$|I^\alpha f(x)| \leq Ct^\alpha Mf(x) + Ct^{\alpha - \frac{n-\lambda}{p}} |x|^{-\frac{\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}.$$

Minimizing with respect to $t = \left[(Mf(x))^{-1} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}} \right]^{\frac{p}{n-\lambda}} |x|^{-\frac{\gamma}{n-\lambda}}$ we arrive at

$$|I^\alpha f(x)| \leq C \left(\frac{Mf(x)}{\|f\|_{L_{p,\lambda,|\cdot|^\gamma}}} \right)^{1 - \frac{p\alpha}{n-\lambda}} |x|^{-\frac{\gamma\alpha}{n-\lambda}}.$$

It is obvious that

$$|x|^\gamma = |x|^{\mu - \frac{\gamma\alpha q}{n-\lambda}}.$$

From Theorem 3.2, taking $\varphi(x) = |x|^\gamma$ we get

$$\begin{aligned} \int_{B(x,t)} |I^\alpha f(y)|^q |y|^\mu dy &\leq C \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}^{q-p} \int_{B(x,t)} (Mf(y))^p |y|^\gamma dy \\ &\leq Ct^\lambda \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}^{q-p} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}^p \\ &= Ct^\lambda \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}^q. \end{aligned}$$

Therefore $I^\alpha f \in L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$ and we obtain

$$\|I^\alpha f\|_{L_{q,\lambda,|\cdot|^\mu}} \leq C \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}.$$

Necessity: Let I^α be bounded from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$, $1 < p < \frac{n-\lambda}{\alpha}$. Define $f_t(x) =: f(tx)$, $t > 0$. Then

$$\begin{aligned} \left(r^{-\lambda} \int_{B(x,r)} |f_t(y)|^p |y|^\gamma dy \right)^{1/p} &= t^{-\frac{n+\gamma}{p}} \left(r^{-\lambda} \int_{B(x,tr)} |f(y)|^p |y|^\gamma dy \right)^{1/p} \\ &= t^{-\frac{n-\lambda+\gamma}{p}} \left((tr)^{-\lambda} \int_{B(x,tr)} |f(y)|^p |y|^\gamma dy \right)^{1/p} \\ &\leq t^{-\frac{n-\lambda+\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}. \end{aligned}$$

Therefore we get

$$\|f_t\|_{L_{p,\lambda,|\cdot|^\gamma}} \leq t^{-\frac{n-\lambda+\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}.$$

Since

$$I^\alpha f_t(x) = t^{-\alpha} I^\alpha f(tx),$$

we obtain

$$\begin{aligned} \left(r^{-\lambda} \int_{B(x,r)} |I^\alpha f_t(y)|^q |y|^\mu dy \right)^{1/q} &= t^{-\alpha} \left(r^{-\lambda} \int_{B(x,r)} |I^\alpha f(ty)|^q |y|^\mu dy \right)^{1/q} \\ &= t^{-\alpha - \frac{n-\lambda+\mu}{q}} \left((tr)^{-\lambda} \int_{B(x,tr)} |I^\alpha f(y)|^q |y|^\mu dy \right)^{1/q} \\ &\leq t^{-\alpha - \frac{n-\lambda+\mu}{q}} \|I^\alpha f\|_{L_{q,\lambda,|\cdot|^\mu}}. \end{aligned}$$

Therefore we get

$$\|I^\alpha f_t\|_{L_{q,\lambda,|\cdot|^\mu}} \leq t^{-\alpha - \frac{n-\lambda+\mu}{q}} \|I^\alpha f\|_{L_{q,\lambda,|\cdot|^\mu}}.$$

Since the operator I^α is bounded from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$, we have

$$\|I^\alpha f_t\|_{L_{q,\lambda,|\cdot|^\mu}} \leq Ct^{-\alpha - \frac{n-\lambda+\mu}{q} + \frac{n-\lambda+\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}, \tag{3.3}$$

where C depends on $p, q, \lambda, \gamma, \mu$ and n .

If $\frac{1}{p} > \frac{1}{q} + \frac{\alpha}{n-\lambda}$, from the inequality (3.3), $\|I^\alpha f_t\|_{L_{q,\lambda,|\cdot|^\mu}} = 0$ for all $f \in L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ as $t \rightarrow 0$.

If $\frac{1}{p} < \frac{1}{q} + \frac{\alpha}{n-\lambda}$, from the inequality (3.3), $\|I^\alpha f_t\|_{L_{q,\lambda,|\cdot|^\mu}} = 0$ for all $f \in L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ as $t \rightarrow \infty$. Therefore $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$. \square

Remark 3.4. The proof of the sufficiency part of Theorem 3.3 is also given with different methods in [26].

Corollary 3.5. [26] *Let $0 < \alpha < n$, $0 \leq \lambda < n - \alpha$, $1 < p < \frac{n-\lambda}{\alpha}$, $-n + \lambda \leq \gamma < n(p - 1) + \lambda$, $\mu = \frac{q\gamma}{p}$ and $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$. Then the operator M^α is bounded from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$.*

4. Commutators of the Riesz potential operator in the spaces

$$L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$$

It is well-known that the commutator is an important integral operator and it plays a key role in harmonic analysis. In this section we consider commutators of the Riesz potential defined by the following equality

$$[b, I^\alpha]f(x) = \int_{\mathbb{R}^n} (b(x) - b(y)) |x - y|^{\alpha-n} f(y) dy, \quad 0 < \alpha < n.$$

Given a measurable function b the operator $|b, I^\alpha|$ is defined by

$$|b, I^\alpha|f(x) = \int_{\mathbb{R}^n} |b(x) - b(y)| |x - y|^{\alpha-n} |f(y)| dy, \quad 0 < \alpha < n.$$

The following statement holds:

Lemma 4.1. [9] *Let $1 < s < \infty$ and $b \in BMO(\mathbb{R}^n)$. Then there exists a positive constant C , independent of f and x , such that*

$$M^\sharp([b, I^\alpha]f(x)) \leq C \|b\|_{BMO} \left[(M|I^\alpha f(x)|^s)^{\frac{1}{s}} + (M^{s\alpha}|f(x)|^s)^{\frac{1}{s}} \right].$$

Proposition 4.2. ([36], Lemma 3.5) *Let $1 < p < \infty$. Then for all $f \in L^p(\mathbb{R}^n)$ and $g \in L^{p'}(\mathbb{R}^n)$ there exists a positive constant C such that*

$$\left| \int_{\mathbb{R}^n} f(y)g(y)dy \right| \leq C \left| \int_{\mathbb{R}^n} M^\sharp f(y)Mg(y)dy \right|.$$

The following lemma is valid.

Lemma 4.3. *Let $1 < p < \infty$, $\varphi \in A_p(\mathbb{R}^n)$. Then there exists a positive constant C , independent of f , such that*

$$\|f\varphi^{\frac{1}{p}}\|_{L_p(\mathbb{R}^n)} \leq C \|\varphi^{\frac{1}{p}}M^\sharp f\|_{L_p(\mathbb{R}^n)}.$$

Proof. By (2.1) we have

$$\|f\varphi^{\frac{1}{p}}\|_{L_p(\mathbb{R}^n)} \leq C \sup_{\|g\|_{L_{p'}(\mathbb{R}^n)} \leq 1} \left| \int_{\mathbb{R}^n} f(y)g(y)\varphi^{\frac{1}{p}}(y)dy \right|.$$

According to Proposition 4.2,

$$\|f\varphi^{\frac{1}{p}}\|_{L_p(\mathbb{R}^n)} \leq C \sup_{\|g\|_{L_{p'}(\mathbb{R}^n)} \leq 1} \left| \int_{\mathbb{R}^n} M^\sharp f(y)M(g\varphi^{\frac{1}{p}})(y)dy \right|.$$

From Hölder inequality and Theorem 3.1, we obtain

$$\begin{aligned} \|f\varphi^{\frac{1}{p}}\|_{L_p(\mathbb{R}^n)} &\leq C \sup_{\|g\|_{L_{p'}(\mathbb{R}^n)} \leq 1} \|\varphi^{\frac{1}{p}}M^\sharp f\|_{L_p(\mathbb{R}^n)} \|\varphi^{-\frac{1}{p}}M(g\varphi^{\frac{1}{p}})\|_{L_{p'}(\mathbb{R}^n)} \\ &\leq C \sup_{\|g\|_{L_{p'}(\mathbb{R}^n)} \leq 1} \|\varphi^{\frac{1}{p}}M^\sharp f\|_{L_p(\mathbb{R}^n)} \|g\|_{L_{p'}(\mathbb{R}^n)} \leq C \|\varphi^{\frac{1}{p}}M^\sharp f\|_{L_p(\mathbb{R}^n)}. \quad \square \end{aligned}$$

Corollary 4.4. *Let $1 < p < \infty$, $\varphi = \psi|\cdot|^\gamma \in A_p(\mathbb{R}^n)$. Then there exists a positive constant C , independent of f , such that*

$$\|f\psi^{\frac{1}{p}}\|_{L_{p,|\cdot|^\gamma}(\mathbb{R}^n)} \leq C \|\psi^{\frac{1}{p}}M^\sharp f\|_{L_{p,|\cdot|^\gamma}(\mathbb{R}^n)}.$$

Lemma 4.5. *Let $1 < p < \infty$, $0 \leq \lambda < n$. Then the following inequality holds*

$$\|f\|_{L_{p,\lambda,|\cdot|^\gamma}} \leq C \|M^\sharp f\|_{L_{p,\lambda,|\cdot|^\gamma}}.$$

Proof. If $0 < \theta < 1$, $\psi(x) = (M\chi_{B(x,r)})^\theta \in A_p(\mathbb{R}^n)$, from Lemma 4.3 we have

$$\|f\|_{L_{p,|\cdot|^\gamma}(B(x,r))} \leq \|f\psi^{\frac{1}{p}}\|_{L_{p,|\cdot|^\gamma}(\mathbb{R}^n)} \leq C \|\psi^{\frac{1}{p}}M^\sharp f\|_{L_{p,|\cdot|^\gamma}(\mathbb{R}^n)} \leq C \|M^\sharp f\|_{L_{p,|\cdot|^\gamma}(B(x,r))}.$$

Therefore we get

$$\begin{aligned} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}} &= \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{L_{p,|\cdot|^\gamma}(B(x,t))} \\ &\leq C \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|M^\sharp f\|_{L_{p,|\cdot|^\gamma}(B(x,r))} = C \|M^\sharp f\|_{L_{p,\lambda,|\cdot|^\gamma}}. \end{aligned}$$

Thus the lemma has been proved. □

In the following theorem we give the necessary and sufficient conditions for the boundedness of the commutator $[b, I^\alpha]$ from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$.

Theorem 4.6. *Let $0 < \alpha < n$, $0 \leq \lambda < n - \alpha$, $1 < p < \frac{n-\lambda}{\alpha}$, $-n + \lambda \leq \gamma < n(p-1) + \lambda$, $\mu = \frac{q\gamma}{p}$ and $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$. Then the commutator $[b, I^\alpha]$ is bounded from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$ if and only if $b \in BMO$.*

Proof. Let $f \in L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ and $b \in BMO(\mathbb{R}^n)$. From Lemma 4.5, we have

$$\|[b, I^\alpha]f\|_{L_{q,\lambda,|\cdot|^\mu}} \leq C_1 \|M^\sharp([b, I^\alpha]f)\|_{L_{q,\lambda,|\cdot|^\mu}}.$$

From Lemma 4.1, we get

$$\begin{aligned} \|M^\sharp([b, I^\alpha]f)\|_{L_{q,\lambda,|\cdot|^\mu}} &\leq C_2 \|b\|_{BMO} \left\| (M|I^\alpha f|^s)^{\frac{1}{s}} + (M^{\alpha s}|f|^s)^{\frac{1}{s}} \right\|_{L_{q,\lambda,|\cdot|^\mu}} \\ &\leq C_3 \|b\|_{BMO} \left[\left\| (M|I^\alpha f|^s)^{\frac{1}{s}} \right\|_{L_{q,\lambda,|\cdot|^\mu}} + \left\| (M^{\alpha s}|f|^s)^{\frac{1}{s}} \right\|_{L_{q,\lambda,|\cdot|^\mu}} \right]. \end{aligned}$$

From Theorem 3.2 and Theorem 3.3, we have

$$\begin{aligned} \left\| (M|I^\alpha f|^s)^{\frac{1}{s}} \right\|_{L_{q,\lambda,|\cdot|^\mu}} &= \|M|I^\alpha f|^s\|_{L_{\frac{q}{s},\lambda,|\cdot|^\mu}}^{\frac{1}{s}} \\ &\leq C \| |I^\alpha f|^s \|_{L_{\frac{q}{s},\lambda,|\cdot|^\mu}}^{\frac{1}{s}} = C \|I^\alpha f\|_{L_{q,\lambda,|\cdot|^\mu}} \leq C \|f\|_{L_{p,\lambda,|\cdot|^\mu}}. \end{aligned}$$

Similarly it can be shown that

$$\left\| (M^{\alpha s}|f|^s)^{\frac{1}{s}} \right\|_{L_{q,\lambda,|\cdot|^\mu}} \leq C \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}.$$

Therefore we obtain

$$\|[b, I^\alpha]f\|_{L_{q,\lambda,|\cdot|^\mu}} \leq C_2 \|b\|_{BMO} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}.$$

(i) \Rightarrow (ii) Now, let us prove the "only if" part. Let $[b, I^\alpha]$ be bounded from $L_{p,\lambda,|\cdot|^\gamma}$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$, $1 < p < \frac{n-\lambda}{\alpha}$. Now we consider $f = \chi_{B(x,r)}$. It is easy to compute that

$$\begin{aligned} \|\chi_{B(x,r)}\|_{L_{p,\lambda,|\cdot|^\gamma}} &\approx \sup_{t>0, x \in \mathbb{R}^n} \left(t^{-\lambda} \int_{B(y,t)} \chi_{B(x,r)}(y) |y|^\gamma dy \right)^{1/p} \\ &\approx \sup_{B(y,t) \subset B(x,r)} \left(t^{-\lambda} \int_{B(y,t)} |y|^\gamma dy \right)^{1/p} \approx r^{\frac{n-\lambda+\gamma}{p}}. \end{aligned}$$

Then

$$\begin{aligned}
 & \frac{1}{|B(x,t)|} \int_{B(x,t)} |b(z) - b_{B(x,t)}| dz \\
 &= \frac{1}{|B(x,t)|} \int_{B(x,t)} \left| b(z) - \frac{1}{|B(x,t)|} \int_{B(x,t)} b(y) dy \right| dz \\
 &\leq \frac{1}{|B(x,t)|^{1+\frac{\alpha}{n}}} \int_{B(x,t)} \frac{1}{|B(x,t)|^{1-\frac{\alpha}{n}}} \left| \int_{B(x,t)} (b(z) - b(y)) dy \right| dz \\
 &\leq \frac{1}{|B(x,t)|^{1+\frac{\alpha}{n}}} \int_{B(x,t)} \left| \int_{B(x,t)} (b(z) - b(y)) |x - y|^{\alpha-n} dy \right| dz \\
 &\leq \frac{1}{|B(x,t)|^{1+\frac{\alpha}{n}}} \int_{B(x,t)} |[b, I^\alpha] \chi_{B(x,t)}(z)| dz \\
 &\leq C t^{-n-\alpha+\lambda} \|[b, I^\alpha] \chi_{B(x,t)}\|_{L_{q,\lambda,|\cdot|^\mu}} \|\chi_{B(x,t)}\|_{L_{q',\lambda,|\cdot|^{1-\frac{\mu}{q}}}} \\
 &\leq C t^{-n-\alpha+\frac{n-\lambda+\gamma}{p}+n-\frac{n-\lambda+\mu}{q}} \leq C.
 \end{aligned}$$

Hence we get

$$|B(x,t)|^{-1} \int_{B(x,t)} |b(y) - b_{B(x,t)}| dy \leq C,$$

which shows that $b \in BMO(\mathbb{R}^n)$.

Thus the theorem has been proved. □

Theorem 4.7. *Let $0 < \alpha < n$, $0 \leq \lambda < n - \alpha$, $1 < p < \frac{n-\lambda}{\alpha}$, $-n + \lambda \leq \gamma < n(p-1) + \lambda$, $\mu = \frac{q\gamma}{p}$ and $b \in BMO$. Then the commutator $|b, I^\alpha|$ is bounded from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$ if and only if $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$.*

Proof. 1) The sufficiency follows from Theorem 4.6.

Necessity: Let $1 < p < \frac{n-\lambda}{\alpha}$ and $|b, I^\alpha|$ be bounded from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$. Define $f_t(x) =: f(tx)$, $t > 0$. Then

$$\begin{aligned}
 \left(r^{-\lambda} \int_{B(x,r)} |f_t(y)|^p |y|^\gamma dy \right)^{1/p} &= t^{-\frac{n+\gamma}{p}} \left(r^{-\lambda} \int_{B(x,tr)} |f(y)|^p |y|^\gamma dy \right)^{1/p} \\
 &= t^{-\frac{n-\lambda+\gamma}{p}} \left((tr)^{-\lambda} \int_{B(x,tr)} |f(y)|^p |y|^\gamma dy \right)^{1/p} \\
 &\leq t^{-\frac{n-\lambda+\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}.
 \end{aligned}$$

Therefore we get

$$\|f_t\|_{L_{p,\lambda,|\cdot|^\gamma}} \leq t^{-\frac{n-\lambda+\gamma}{p}} \|f\|_{L_{p,\lambda,|\cdot|^\gamma}}.$$

Since

$$|b, I^\alpha|f_t(x) = t^{-\alpha}|b, I^\alpha|f(tx),$$

we obtain

$$\begin{aligned} & \left(r^{-\lambda} \int_{B(x,r)} [||b, I^\alpha|f_t||^q(y)|y|^\mu dy] \right)^{1/q} \\ &= t^{-\alpha} \left(r^{-\lambda} \int_{B(x,r)} [||b, I^\alpha|f||^q(ty)|y|^\mu dy] \right)^{1/q} \\ &= t^{-\alpha - \frac{n-\lambda+\mu}{q}} \left((tr)^{-\lambda} \int_{B(x,tr)} [||b, I^\alpha|f||^q(y)|y|^\mu dy] \right)^{1/q} \\ &\leq t^{-\alpha - \frac{n-\lambda+\mu}{q}} ||b, I^\alpha|f||_{L_{q,\lambda,|\cdot|^\mu}}. \end{aligned}$$

Therefore we get

$$||b, I^\alpha|f_t||_{L_{q,\lambda,|\cdot|^\mu}} \leq t^{-\alpha - \frac{n-\lambda+\mu}{q}} ||b, I^\alpha|f||_{L_{q,\lambda,|\cdot|^\mu}}.$$

Since the operator $|b, I^\alpha|$ is bounded from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$, we have

$$||b, I^\alpha|f_t||_{L_{q,\lambda,|\cdot|^\mu}} \leq C t^{-\alpha - \frac{n-\lambda+\mu}{q} + \frac{n-\lambda+\gamma}{p}} ||b||_{BMO} ||f||_{L_{p,\lambda,|\cdot|^\gamma}}, \tag{4.1}$$

where C depends on $p, q, \lambda, \gamma, \mu$ and n .

If $\frac{1}{p} > \frac{1}{q} + \frac{\alpha}{n-\lambda}$, from the inequality (4.1), $||b, I^\alpha|f_t||_{L_{q,\lambda,|\cdot|^\mu}} = 0$ for all $f \in L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ as $t \rightarrow 0$.

If $\frac{1}{p} < \frac{1}{q} + \frac{\alpha}{n-\lambda}$, from the inequality (4.1), $||b, I^\alpha|f_t||_{L_{q,\lambda,|\cdot|^\mu}} = 0$ for all $f \in L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ as $t \rightarrow \infty$. Therefore $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$. □

The following theorem gives the conditions for the boundedness of the commutator $|b, I^\alpha|$ from $B_{p,\theta,\lambda,|\cdot|^\gamma}^s(\mathbb{R}^n)$ to $B_{q,\theta,\lambda,|\cdot|^\mu}^s(\mathbb{R}^n)$.

Theorem 4.8. *Let $0 < \alpha < n$, $0 \leq \lambda < n - \alpha$, $1 < p < \frac{n-\lambda}{\alpha}$, $-n + \lambda \leq \gamma < n(p-1) + \lambda$, $\mu = \frac{q\gamma}{p}$, $0 < s < 1$, $1 \leq \theta \leq \infty$, $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$ and $b \in BMO(\mathbb{R}^n)$. Then the commutator $|b, I^\alpha|$ is bounded from $B_{p,\theta,\lambda,|\cdot|^\gamma}^s(\mathbb{R}^n)$ to $B_{q,\theta,\lambda,|\cdot|^\mu}^s(\mathbb{R}^n)$.*

Proof. From the definition of the Besov-Morrey type spaces it suffices to show that

$$||b, I^\alpha|f(x - \cdot) - |b, I^\alpha|f(\cdot)||_{L_{p,\lambda,|\cdot|^\gamma}} \leq C ||b||_{BMO} ||f(x - \cdot) - f(\cdot)||_{L_{p,\lambda,|\cdot|^\gamma}}.$$

Hence we have

$$|[b, I^\alpha]f(x - \cdot) - |b, I^\alpha|f| \leq |b, I^\alpha|(|f(x - \cdot) - f|).$$

Taking $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ norm of both sides of the above inequality, from the boundedness of $|b, I^\alpha|$ from $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|^\mu}(\mathbb{R}^n)$, we obtain the desired result. Thus Theorem 4.8 has been proved. □

5. The weighted Morrey estimates for the operators $V^s(-\Delta + V)^{-\beta}$ and $V^s\nabla(-\Delta + V)^{-\beta}$

In this section we consider the Schrödinger operator $-\Delta + V$ on \mathbb{R}^n , where the nonnegative potential V belongs to the reverse Hölder class $B_q(\mathbb{R}^n)$ for some $q_1 \geq n$. We obtain weighted Morrey $L_{p,\lambda,|\cdot|^\gamma}(\mathbb{R}^n)$ estimates for the operators $V^s(-\Delta + V)^{-\beta}$ and $V^s\nabla(-\Delta + V)^{-\beta}$.

Schrödinger operators on the Euclidean space \mathbb{R}^n with nonnegative potentials which belong to the reverse Hölder class have been studied by many authors (see [10, 32, 40]). Shen [32] studied the Schrödinger operator $-\Delta + V$, assuming the nonnegative potential V belongs to the reverse Hölder class $B_q(\mathbb{R}^n)$ for $q \geq n/2$ and he proved the L_p boundedness of the operators $(-\Delta + V)^{is}$, $\nabla^2(-\Delta + V)^{-1}$, $\nabla(-\Delta + V)^{-\frac{1}{2}}$ and $\nabla(-\Delta + V)^{-1}$. Kurata and Sugano generalized Shens' results to uniformly elliptic operators in [18]. Sugano [38] also extended some results of Shen to the operator $V^s(-\Delta + V)^{-\beta}$, $0 \leq s \leq \beta \leq 1$ and $V^s\nabla(-\Delta + V)^{-\beta}$, $0 \leq s \leq \frac{1}{2} \leq \beta \leq 1$ and $\beta - s \geq \frac{1}{2}$. Later, Lu [21] and Li [19] investigated the Schrödinger operators in a more general setting.

We investigate the weighted Morrey $L_{p,\lambda,|\cdot|^\gamma} - L_{q,\lambda,|\cdot|^\mu}$ boundedness of the operators

$$T_1 = V^s(-\Delta + V)^{-\beta}, \quad 0 \leq s \leq \beta \leq 1,$$

$$T_2 = V^s\nabla(-\Delta + V)^{-\beta}, \quad 0 \leq s \leq \frac{1}{2} \leq \beta \leq 1, \quad \beta - s \geq \frac{1}{2}.$$

Note that the operators $V(-\Delta + V)^{-1}$ and $V^{\frac{1}{2}}\nabla(-\Delta + V)^{-1}$ in [19] are the special case of T_1 and T_2 , respectively.

It is worth pointing out that we need to establish pointwise estimates for T_1 , T_2 and their adjoint operators by using the estimates of fundamental solution for the Schrödinger operator on \mathbb{R}^n in [19]. And we give the Morrey estimates by using $L_{p,\lambda,|\cdot|^\gamma} - L_{q,\lambda,|\cdot|^\mu}$ boundedness of the fractional maximal operators.

Definition 5.1. 1) A nonnegative locally L_p integrable function V on \mathbb{R}^n is said to belong to the reverse Hölder class B_p ($1 < p < \infty$) if there exists a positive constant C such that the reverse Hölder inequality

$$\left(\frac{1}{|B|} \int_B V(x)^p dx \right)^{\frac{1}{p}} \leq \frac{C}{|B|} \int_B V(x) dx$$

holds for every ball B in \mathbb{R}^n .

2) Let $V \geq 0$. We say $V \in B_\infty$, if there exists a positive constant C such that the inequality

$$\|V\|_{L_\infty(B)} \leq \frac{C}{|B|} \int_B V(x) dx$$

holds for every ball B in \mathbb{R}^n .

Clearly, $B_\infty \subset B_p$ for $1 < p < \infty$. But it is important that the B_p class has a property of "self-improvement"; that is, if $V \in B_p$, then $V \in B_{p+\varepsilon}$ for some $\varepsilon > 0$ (see [19]).

The following two pointwise estimates for T_1 and T_2 were proved in [40] with the potential $V \in B_\infty$.

Theorem A. *Suppose $V \in B_\infty$ and $0 \leq s \leq \beta \leq 1$. Then there exists a positive constant C such that*

$$|T_1 f(x)| \leq CM^\alpha f(x), \quad f \in C_0^\infty(\mathbb{R}^n),$$

where $\alpha = 2(\beta - s)$.

Theorem B. *Suppose $V \in B_\infty$, $0 \leq s \leq \frac{1}{2} \leq \beta \leq 1$ and $\beta - s \geq \frac{1}{2}$. Then there exists a positive constant C such that*

$$|T_2 f(x)| \leq CM^\alpha f(x), \quad f \in C_0^\infty(\mathbb{R}^n),$$

where $\alpha = 2(\beta - s) - 1$.

Note that the similar estimates for the adjoint operators T_1^* and T_2^* with the potential $V \in B_{q_1}$ for some $q_1 > \frac{n}{2}$ are also valid (see [20]).

Theorem C. *Suppose $V \in B_{q_1}$ for some $q_1 > \frac{n}{2}$, $0 \leq s \leq \beta \leq 1$ and let $\frac{1}{q_2} = 1 - \frac{\alpha}{q_1}$. Then there exists a positive constant C such that*

$$|T_1^* f(x)| \leq C (M_{\alpha q_2}(|f|^{q_2})(x))^{\frac{1}{q_2}}, \quad f \in C_0^\infty(\mathbb{R}^n),$$

where $\alpha = 2(\beta - s)$.

Theorem D. *Suppose $V \in B_{q_1}$ for some $q_1 > \frac{n}{2}$, $0 \leq s \leq \frac{1}{2} \leq \beta \leq 1$ and $\beta - s \geq \frac{1}{2}$. And let*

$$\frac{1}{q_1} = \begin{cases} 1 - \frac{s}{q_1}, & \text{if } q_1 > n, \\ 1 - \frac{\alpha+1}{q_1} + \frac{1}{n}, & \text{if } \frac{n}{2} < q_1 < n. \end{cases}$$

Then there exists a positive constant C such that

$$|T_2^* f(x)| \leq C (M_{\alpha q_2}(|f|^{q_2})(x))^{\frac{1}{q_2}}, \quad f \in C_0^\infty(\mathbb{R}^n),$$

where $\alpha = 2(\beta - s) - 1$.

The above theorems will yield the weighted Morrey estimates for T_1 and T_2 .

Corollary 5.2. *Assume that $V \in B_\infty$, and $0 \leq s \leq \beta \leq 1$. Let $1 < p < \frac{n}{s}$, $-n + \lambda \leq \gamma < n(p - 1) + \lambda$, $\mu = \frac{q\gamma}{p}$, $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$ and $0 \leq \lambda < n$, where $\alpha = 2(\beta - s) < n$.*

Then for any $f \in C_0^\infty(\mathbb{R}^n)$ there exists a positive constant C such that

$$\|T_1 f\|_{L_{q,\lambda,|\cdot|}^\mu} \leq C \|f\|_{L_{p,\lambda,|\cdot|}^\gamma}.$$

Corollary 5.3. *Let $V \in B_\infty$, $0 \leq s \leq \frac{1}{2} \leq \beta \leq 1$, $\beta - s \geq \frac{1}{2}$, $1 < p < \frac{n}{\alpha}$, $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$, $-n + \lambda \leq \gamma < n(p - 1) + \lambda$, $\mu = \frac{q\gamma}{p}$ and $0 \leq \lambda < n$, where $\alpha = 2(\beta - s) - 1 < n$.*

Then for any $f \in C_0^\infty(\mathbb{R}^n)$ there exists a positive constant C such that

$$\|T_2 f\|_{L_{q,\lambda,|\cdot|}^\mu} \leq C \|f\|_{L_{p,\lambda,|\cdot|}^\gamma}.$$

Corollary 5.4. *Assume that $V \in B_{q_1}$ for $q_1 > \frac{n}{2}$, and $0 \leq s \leq \beta \leq 1$.*

Let $\frac{1}{q_2} = 1 - \frac{\alpha}{q_1}$, $1 < p < \frac{1}{\frac{\alpha}{q_1} + \frac{1}{n}}$, $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{\frac{n}{q_2} - \lambda}$, $-n + \lambda \leq \gamma < n(p - 1) + \lambda$, $\mu = \frac{q\gamma}{p}$ and $0 \leq \lambda < nq_2$, where $\alpha = 2(\beta - s) < n$.

Then for any $f \in C_0^\infty(\mathbb{R}^n)$ there exists a positive constant C such that

$$\|T_1 f\|_{L_{q,\lambda,|\cdot|}^\mu} \leq C \|f\|_{L_{p,\lambda,|\cdot|}^\gamma}.$$

Corollary 5.5. *Assume that $V \in B_{q_1}$ for $q_1 > \frac{n}{2}$, and*

$$\begin{cases} 0 \leq s \leq \frac{1}{2} \leq \beta \leq 1, & \text{if } q_1 > n, \\ 0 \leq s \leq \frac{1}{2} < \beta \leq 1, & \text{if } \frac{n}{2} < q_1 < n. \end{cases}$$

Let $\alpha = 2(\beta - s) - 1 < n$ and $\beta - s \geq \frac{1}{2}$, and let $1 < p < \frac{1}{\frac{\alpha}{q_1} + \frac{\alpha}{n}}$, $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{\frac{n}{q_2} - \lambda}$, $\frac{1}{q_2} = 1 - \frac{\alpha}{q_1}$, $-n + \lambda \leq \gamma < n(p - 1) + \lambda$, $\mu = \frac{q\gamma}{p}$ and $0 \leq \lambda < nq_2$, where

$$\frac{1}{p_1} = \begin{cases} \frac{\alpha}{q_1}, & \text{if } q_1 > n, \\ \frac{\alpha+1}{q_1} + \frac{1}{n}, & \text{if } \frac{n}{2} < q_1 < n. \end{cases}$$

Then for any $f \in C_0^\infty(\mathbb{R}^n)$ there exists a positive constant C such that

$$\|T_2 f\|_{L_{q,\lambda,|\cdot|}^\mu} \leq C \|f\|_{L_{p,\lambda,|\cdot|}^\gamma}.$$

6. Some applications

The theorems of the Section 3 can be applied to various operators which are estimated from above by Riesz potentials. Now we give some examples.

Suppose that L is a linear operator on L_2 which generates an analytic semigroup e^{-tL} with the kernel $p_t(x, y)$ satisfying a Gaussian upper bound, that is,

$$|p_t(x, y)| \leq \frac{c_1}{t^{n/2}} e^{-c_2 \frac{|x-y|^2}{t}} \tag{6.1}$$

for $x, y \in \mathbb{R}^n$ and all $t > 0$.

For $0 < \alpha < n$, the fractional powers $L^{-\alpha/2}$ of the operator L are defined by

$$L^{-\alpha/2} f(x) = \frac{1}{\Gamma(\alpha/2)} \int_0^\infty e^{-tL} f(x) \frac{dt}{t^{-\alpha/2+1}}.$$

Note that if $L = -\Delta$ is the Laplacian on \mathbb{R}^n , then $L^{-\alpha/2}$ is the Riesz potential I^α . (See, for example, Chapter 5 in [36].)

Theorem 6.1. *Let $0 < \alpha < n$, $0 \leq \lambda < n - \alpha$, $1 < p < \frac{n-\lambda}{\alpha}$, $-n + \lambda \leq \gamma < n(p - 1) + \lambda$, $\mu = \frac{q\gamma}{p}$ and condition (6.1) be satisfied. Then condition $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$ is sufficient for the boundedness of $L^{-\alpha/2}$ from $L_{p,\lambda,|\cdot|}^\gamma(\mathbb{R}^n)$ to $L_{q,\lambda,|\cdot|}^\mu(\mathbb{R}^n)$.*

Proof. Since the semigroup e^{-tL} has the kernel $p_t(x, y)$ which satisfies condition (6.1), it follows that

$$|L^{-\alpha/2} f(x)| \leq C I^\alpha |f|(x)$$

for all $x \in \mathbb{R}^n$ (see [7]). Therefore from the aforementioned theorems we have

$$\|L^{-\alpha/2} f\|_{L_{q,\lambda,|\cdot|}^\mu} \leq C \|I^\alpha |f|\|_{L_{q,\lambda,|\cdot|}^\mu} \leq C \|f\|_{L_{p,\lambda,|\cdot|}^\gamma}. \quad \square$$

Large classes of differential operators satisfies condition (6.1). Now we investigate two of them:

(i) Let us consider a magnetic potential \vec{a} , i. e., a real-valued vector potential $\vec{a} = (a_1, a_2, \dots, a_n)$, and an electric potential V . Assume that for any $k = 1, 2, \dots, n$, $a_k \in L_2^{loc}$ and $0 \leq V \in L_1^{loc}$. The magnetic Schrödinger operator, L , is defined by

$$L = -(\nabla - i\vec{a})^2 + V(x).$$

From the well-known diamagnetic inequality (see [35], Theorem 2.3) we have the following pointwise estimate. For any $t > 0$ and $f \in L_2$,

$$|e^{-tL}f| \leq e^{-t\Delta}|f|,$$

which implies that the semigroup e^{-tL} has the kernel $p_t(x, y)$ that satisfies upper bound (6.1).

(ii) Let $A = (a_{ij}(x))_{1 \leq i, j \leq n}$ be an $n \times n$ matrix with complex-valued entries $a_{ij} \in L_\infty$ satisfying

$$\operatorname{Re} \sum_{i, j=1}^n a_{ij}(x) \zeta_i \zeta_j \geq \lambda |\zeta|^2$$

for all $x \in \mathbb{R}^n$, $\zeta = (\zeta_1, \zeta_2, \dots, \zeta_n) \in \mathbb{C}^n$ and some $\lambda > 0$. Consider the divergence form operator

$$Lf \equiv -\operatorname{div}(A\nabla f),$$

which is interpreted in the usual weak sense via the appropriate sesquilinear form.

It is known that the Gaussian bound (6.1) for the kernel of e^{-tL} holds when A has real-valued entries (see, for example, [3]), or when $n = 1, 2$ in the case of complex-valued entries (see [4, Chapter 1]).

Finally we note that under the appropriate assumptions (see [23]; [36], Chapter 5; [4], pp. 58-59) one can obtain results similar to Theorem 6.1 for a homogeneous elliptic operator L in L_2 of order $2m$ in the divergence form

$$Lf = (-1)^m \sum_{|\alpha|=|\beta|=m} D^\alpha (a_{\alpha\beta} D^\beta f).$$

In this case estimate (6.1) should be replaced by

$$|p_t(x, y)| \leq \frac{c_3}{t^{n/2m}} e^{-c_4 \left(\frac{|x-y|}{t^{1/(2m)}} \right)^{2m/(2m-1)}}$$

for all $t > 0$ and all $x, y \in \mathbb{R}^n$.

References

- [1] Adams, D.R., *A note on Riesz potentials*, Duke Math. J., **42**(1975), 765-778.
- [2] Adams, D.R., *Choquet integrals in potential theory*, Publ. Mat., **42**(1998), 3-66.
- [3] Arendt, W., Elst, A.F.M., *Gaussian estimates for second order elliptic operators with boundary conditions*, J. Operator Theory, **38**(1997), 87-130.
- [4] Auscher, P., Tchamitchian, P., *Square Root Problem for Divergence Operators and Related Topics*, Astérisque, 249, Soc. Math. France, 1998.
- [5] Caffarelli, L., *Elliptic second order equations*, Rend. Sem. Mat. Fis. Milano, **58**(1990), 253-284.
- [6] Chiarenza, F., Frasca, M., *Morrey spaces and Hardy-Littlewood maximal function*, Rend. Math., **7**(1987), 273-279.
- [7] Duong, X.T., Yan, L.X., *On commutators of fractional integrals*, Proc. Amer. Math. Soc., **132**(2004), no. 12, 3549-3557.

- [8] Di Fazio, G., Palagachev, D.K., Ragusa, M.A., *Global Morrey regularity of strong solutions to the Dirichlet problem for elliptic equations with discontinuous coefficients*, J. Funct. Anal., **166**(1999), 179-196.
- [9] Di Fazio, G., Ragusa, M.A., *Commutators and Morrey spaces*, Bollettino U.M.I., **7**(5-A)(1991), 323-332.
- [10] Fefferman, C., *The uncertainty principle*, Bull. Amer. Math. Soc., **9**(1983), 129-206.
- [11] Giaquinta, M., *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, Princeton Univ. Press, Princeton, NJ, 1983.
- [12] Hardy, H.G., Littlewood, J.E., *Some properties of fractional integrals*, I. Math. Z., **27**(1928), no. 4, 565-606.
- [13] Hasanov, J.J., *Hardy-Littlewood-Stein-Weiss inequality in the variable exponent Morrey spaces*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb., **39**(2013), 47-62.
- [14] Ho, K-P., *Singular integral operators, John-Nirenberg inequalities and Tribel-Lizorkin type spaces on weighted Lebesgue spaces with variable exponents*, Rev. Un. Mat. Argentina, **57**(2016), no. 1, 85-101.
- [15] Karaman, T., Guliyev, V.S., Serbetci, A., *Boundedness of sublinear operators generated by Calderón-Zygmund operators on generalized weighted Morrey spaces*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.), **60**(2014), no. 1, 227-244.
- [16] Kufner, A., John, O., Fučík, S., *Function Spaces*, Noordhoff International Publishing: Leyden, Publishing House Czechoslovak Academy of Sciences: Prague, 1977.
- [17] Kurata, K., Nishigaki, S., Sugano, S., *Boundedness of integral operators on generalized Morrey spaces and its application to Schrödinger operators*, Proc. AMS, **128**(1999), no. 4, 1125-1134.
- [18] Kurata, K., Sugano, S., *A remark on estimates for uniformly elliptic operators on weighted L_p spaces and Morrey spaces*, Math. Nachr., **209**(2000), 137-150.
- [19] Li, H.Q., *Estimations L_p des operateurs de Schrödinger sur les groupes nilpotents*, J. Funct. Anal., **161**(1999), 152-218.
- [20] Liu, Y., *The weighted estimates for the operators $V^\alpha(-\Delta_G + V)^{-\beta}$ and $V^\alpha \nabla_G(-\Delta_G + V)^{-\beta}$ on the stratified Lie group \mathbb{G}* , J. Math. Anal. Appl., **349**(2009), 235-244.
- [21] Lu, G.Z., *A Fefferman-Phong type inequality for degenerate vector fields and applications*, Panamer. Math. J., **6**(1996), 37-57.
- [22] Mazzucato, A.L., *Besov-Morrey spaces: Function space theory and applications to non-linear PDE*, Trans. Amer. Math. Soc., **355**(2003), 1297-1364.
- [23] McIntosh, A., *Operators which have an H_1 -calculus*, in Proc. Centre Math. Analysis, Vol. 14, Miniconference on Operator Theory and Partial Differential Equations, A.N.U., Canberra, 1986, 210-231.
- [24] Morrey, C.B., *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc., **43**(1938), 126-166.
- [25] Muckenhoupt, B., *Weighted norm inequalities for the Hardy-Littlewood maximal function*, Trans. Amer. Math. Soc., **165**(1972), 207-226.
- [26] Nakamura, S., Sawano, Y., Tanaka, H., *The fractional operators on weighted Morrey spaces*, J. Geom. Anal., **28**(2018), no. 2, 1502-1524.
- [27] Peetre, J., *On the theory of $\mathcal{L}^{p,\lambda}$ spaces*, J. Funct. Anal., **4**(1969), 71-87.
- [28] Perez, C., *Two weighted norm inequalities for Riesz potentials and uniform L_p -weighted Sobolev inequalities*, Indiana Univ. Math. J., **39**(1990), 31-44.

- [29] Ruiz, A., Vega, L., *Unique continuation for Schrödinger operators with potential in Morrey spaces*, Publ. Mat., **35**(1991), 291-298.
- [30] Ruiz, A., Vega, L., *On local regularity of Schrödinger equations*, Int. Math. Res. Notices, **1993**(1993), no. 1, 13-27.
- [31] Samko, S., *Hardy-Littlewood-Stein-Weiss inequality in the Lebesgue spaces with variable exponent*, Fract. Calc. Appl. Anal., **6**(2003), no. 4, 421-440.
- [32] Shen, Z., *L_p estimates for Schrödinger operators with certain potentials*, Ann. Inst. Fourier (Grenoble), **45**(1995), 513-546.
- [33] Shen, Z., *The periodic Schrödinger operators with potentials in the Morrey class*, J. Funct. Anal., **193**(2002), 314-345.
- [34] Shen, Z., *Boundary value problems in Morrey spaces for elliptic systems on Lipschitz domains*, Amer. J. Math., **125**(2003), 1079-1115.
- [35] Simon, B., *Maximal and minimal Schrödinger forms*, J. Op. Theory, **1**(1979), 37-47.
- [36] Stein, E.M., *Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton Univ. Press, Princeton, NJ, 1993.
- [37] Stein, E.M., Weiss, G., *Fractional integrals on n -dimensional Euclidean space*, J. Math. and Mech., **7**(1958), no. 4, 503-514.
- [38] Sugano, S., *Estimates for the operators $V^\alpha(-\Delta + V)^{-\beta}$ and $V^\alpha \nabla(-\Delta + V)^{-\beta}$ with certain nonnegative potentials V* , Tokyo J. Math., **21**(1998), 441-452.
- [39] Taylor, M.E., *Analysis on Morrey spaces and applications to Navier-Stokes and other evolution equations*, Comm. Partial Differential Equations, **17**(1992), 1407-1456.
- [40] Zhong, J.P., *Harmonic analysis for some Schrödinger type operators*, PhD Thesis, Princeton University, 1993.

Canay Aykol
Ankara University,
Department of Mathematics
Ankara, Turkey
e-mail: aykol@science.ankara.edu.tr

Javanshir J. Hasanov
Azerbaijan State Oil and Industry,
Baku, Azerbaijan
e-mail: hasanovjavanshir@yahoo.com.tr