

# The Nehari–Schwarz lemma and infinitesimal boundary rigidity of bounded holomorphic functions

Oliver Roth

*Dedicated to the memory of Professor Gabriela Kohr*

**Abstract.** We survey a number of recent generalizations and sharpenings of Nehari’s extension of Schwarz’ lemma for holomorphic self–maps of the unit disk. In particular, we discuss the case of infinitely many critical points and its relation to the zero sets and invariant subspaces for Bergman spaces, as well as the case of equality at the boundary.

**Mathematics Subject Classification (2010):** 30C80, 32A36.

**Keywords:** Schwarz’ lemma, Blaschke products, Bergman spaces.

## 1. Introduction

Let  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$  be the open unit disk in the complex plane  $\mathbb{C}$ , and let  $\mathcal{B}$  denote the set of holomorphic functions from  $\mathbb{D}$  into  $\overline{\mathbb{D}}$ . A *finite Blaschke product of degree  $n$*  is a rational function  $B \in \mathcal{B}$  of the form

$$B(z) = \eta \prod_{j=1}^n \frac{z_j - z}{1 - \overline{z_j}z}, \quad |\eta| = 1, \quad (1.1)$$

with zeros  $z_1, \dots, z_n \in \mathbb{D}$ , not necessarily pairwise distinct. Hence the multiplicative building blocks of finite Blaschke products are exactly the elements of the group of conformal automorphisms of  $\mathbb{D}$ ,

$$\text{Aut}(\mathbb{D}) = \left\{ \eta \frac{z_0 - z}{1 - \overline{z_0}z} : |\eta| = 1, z_0 \in \mathbb{D} \right\}.$$

Blaschke products are omnipresent, and occur for instance as fundamental norm-preserving factors in many important classes of holomorphic functions on  $\mathbb{D}$ . We refer

to the recent monograph [19] and the references therein for an state-of-the-art account of the properties and abundant applications of FBP, the set of all finite Blaschke products. In this note, we discuss a number of recent generalizations of Nehari’s celebrated extension [37] of Schwarz’ lemma, a topic which is intrinsically related to FBPs, but which has not been treated in [19].

As point of departure, we note that a geometric-topological way of thinking about (non-constant) FBPs is to view them as proper holomorphic self-maps of  $\mathbb{D}$  or – equivalently – as finite branched coverings of  $\mathbb{D}$ , see [19, Chapter 3]. From this view point, it seems natural to describe a finite Blaschke product  $B$  not in terms of its zeros as in (1.1), but in terms of its *critical points*, that is, the zeros of its first derivative  $B'$ . That this is indeed possible is the content of the following celebrated result of M. Heins [24] (see also [19, Chapter 6], [47] and [48, 8, 53, 45, 30, 31, 41, 49]).

**Theorem 1.1 (Heins 1962).** *Let  $c_1, \dots, c_{n-1}$  be points in  $\mathbb{D}$ . Then there is a Blaschke product  $B$  of degree  $n$  with critical points  $c_1, \dots, c_{n-1}$  in  $\mathbb{D}$  and no others. The Blaschke product  $B$  is unique up to post-composition with an element of  $\text{Aut}(\mathbb{D})$ .*

The Blaschke product  $B$  in Theorem 1.1 can be characterized as the essentially unique extremal function in a sharpened form of the Schwarz–Pick inequality. This fundamental observation [37] is due to Nehari in 1947. In order to state Nehari’s result we need to introduce some notation. We denote by  $\mathcal{C}_f$  the collection of all critical points of a non-constant function  $f \in \mathcal{B}$  counting multiplicities. By slight abuse of language, we call  $\mathcal{C}_f$  the *critical set of  $f$*  and write  $\mathcal{C}_g \subseteq \mathcal{C}_f$  whenever each critical point of a function  $g \in \mathcal{B}$  is also a critical point of  $f \in \mathcal{B}$  of at least the same multiplicity. This is in accordance with standard practices, see [14, §4.1].

**Theorem 1.2 (The Nehari–Schwarz lemma).** *Let  $f \in \mathcal{B}$  and let  $B \in \text{FBP}$  such that  $\mathcal{C}_B \subseteq \mathcal{C}_f$ . Then:*

(i) *(Nehari–Schwarz inequality)*

$$\frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{|B'(z)|}{1 - |B(z)|^2} \quad \text{for all } z \in \mathbb{D}; \tag{1.2}$$

(ii) *(Strong form of the Nehari–Schwarz lemma at an interior point)*

*Equality holds in (1.2) for some point  $z \in \mathbb{D} \setminus \mathcal{C}_B$  if and only if  $f = T \circ B$  for some  $T \in \text{Aut}(\mathbb{D})$ .*

**Remark 1.3 (The Schwarz–Pick lemma).** In Theorem 1.2, one can always take  $B$  as a finite Blaschke product without any critical points, that is, as a conformal automorphism of  $\mathbb{D}$ . In this case, Theorem 1.2 reduces to the standard Schwarz–Pick lemma:

(i) *(Schwarz–Pick inequality)*

$$\frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z \in \mathbb{D}; \tag{1.3}$$

(ii) *(Strong form of the Schwarz–Pick lemma at an interior point)*

*Equality holds in (1.3) for some point  $z \in \mathbb{D}$  if and only if  $f \in \text{Aut}(\mathbb{D})$ .*

The main purpose of this note is to survey some recent sharpenings and extensions of the Nehari–Schwarz lemma. In Section 3 we discuss a generalization of the Nehari–Schwarz lemma which allows for taking into account *infinitely many* critical points instead of only finitely many as in Theorem 1.2. In Section 4 we describe the connections with the specific Bergman space  $A_1^2$ , in particular its zero sets and invariant subspaces. Our presentation is based on recent work of Kraus [28], Dyakonov [15, 16], and Ivrii [26, 27]. In Section 5 we discuss the so-called strong form of the Nehari–Schwarz lemma, that is, the case of equality in the Nehari–Schwarz inequality *at the boundary* which has recently been obtained in [9, Theorem 2.10] as a special case of a general boundary rigidity theorem for conformal pseudometrics. In order to make this paper self-contained we also provide a fairly concise proof of the Nehari–Schwarz inequality (1.2) in Section 2. The proof we give is slightly different from the standard proofs which can be found in [37, Corollary, p. 1037] and [24, Theorem 24.1]. The Nehari–Schwarz lemma has found many further applications, for which we refer to other works such as [6, 22, 34, 35, 45], for instance.

## 2. Proof of the Nehari–Schwarz inequality

We give a proof of the Nehari–Schwarz inequality (1.2) which is based on the observation that a finite Blaschke product  $B$  has the property that

$$\lim_{|z| \rightarrow 1} (1 - |z|^2) \frac{|B'(z)|}{1 - |B(z)|^2} = 1. \quad (2.1)$$

In fact, condition (2.1) *characterizes* finite Blaschke products (Heins [25], see also [33] and [19, Chapter 6.5]). We point out that a simple and direct proof that (2.1) holds for any finite Blaschke product

$$B(z) = \eta \prod_{j=1}^n \frac{z - z_j}{1 - \bar{z}_j z}$$

is possible by making appeal to an identity due to Frostman [18], namely

$$\frac{1 - |B(z)|^2}{1 - |z|^2} = \sum_{k=1}^n \left( \prod_{j=1}^{k-1} \left| \frac{z - z_j}{1 - \bar{z}_j z} \right|^2 \right) \frac{1 - |z_k|^2}{|1 - \bar{z}_k z|^2}, \quad |z| \neq 1, \quad (2.2)$$

and the elementary formula for the logarithmic derivative of  $B$  given by

$$\frac{B'(z)}{B(z)} = \sum_{k=1}^n \frac{1 - |z_k|^2}{(1 - \bar{z}_k z)(z - z_k)}. \quad (2.3)$$

Frostman's identity (2.2) can be easily established by induction, see [19, p. 77]. Clearly, (2.2) and (2.3) immediately imply (2.1).

Using (2.1) we now give a proof of Theorem 1.2 following very closely the standard proof of Ahlfors' lemma [1] with only minor modifications.

*Proof of Theorem 1.2 (i).* Let  $f \in \mathcal{B}$  be non-constant, so  $\mathcal{C}_f$  is a discrete subset of  $\mathbb{D}$ . We consider the auxiliary function

$$u(z) := \log \left( \frac{|f'(z)|}{1 - |f(z)|^2} \frac{1 - |B(z)|^2}{|B'(z)|} \right).$$

Since  $\mathcal{C}_B \subseteq \mathcal{C}_f$  (“including multiplicities”), we see that  $u$  is well-defined and real analytic on  $\mathbb{D} \setminus \mathcal{C}_f$ . For each  $\xi \in \mathcal{C}_f$  the limit

$$\lim_{z \rightarrow \xi} u(z) \in \mathbb{R} \cup \{-\infty\}$$

exists, so  $u$  extends to an upper semicontinuous function on  $\mathbb{D}$  with values in  $\mathbb{R} \cup \{-\infty\}$  which we continue to denote by  $u$ . Now, a straightforward computation reveals

$$\Delta u = 4 \left( \frac{|B'(z)|}{1 - |B(z)|^2} \right)^2 (e^{2u} - 1), \quad z \in \mathbb{D} \setminus \mathcal{C}_f.$$

In particular,

$$u^+ := \max\{u, 0\}$$

is subharmonic in  $\mathbb{D}$ . On the other hand, in view of the Schwarz–Pick inequality,

$$(1 - |z|^2) \frac{|f'(z)|}{1 - |f(z)|^2} \leq 1,$$

we deduce from (2.1) that

$$\limsup_{|z| \rightarrow 1} u(z) \leq 0.$$

Hence  $u^+ \leq 0$  in  $\mathbb{D}$  by the maximum principle. This implies  $u \leq 0$  and completes the proof of (1.2). □

**Remark 2.1 (Strong form of the Nehari–Schwarz lemma at an interior point).** The case of equality for the Nehari–Schwarz inequality (1.2) for some *interior* point  $z \in \mathbb{D} \setminus \mathcal{C}_B$  can be handled in a similar way as the case of equality at some interior point for Ahlfors’ lemma, which has been treated in [24, 40, 36, 12, 33]. We refer to e.g. [33, Remark 2.2 (d)] for the details.

### 3. Infinitely many critical points

We begin with an extension of the theorems of Heins’ (Theorem 1.1) and Nehari–Schwarz (Theorem 1.2) essentially due to Kraus [28].

**Theorem 3.1.** *Let  $\mathcal{C}$  be the critical set of a non-constant function in  $\mathcal{B}$ . Then there is a Blaschke product  $B$  with critical set  $\mathcal{C}$  such that*

$$\frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{|B'(z)|}{1 - |B(z)|^2}$$

for all  $z \in \mathbb{D}$  and any  $f \in \mathcal{B}$  such that  $\mathcal{C}_f \supseteq \mathcal{C}$ . If equality holds at a single point  $z \notin \mathcal{C}$ , then  $f = T \circ B$  for some  $T \in \text{Aut}(\mathbb{D})$ . The Blaschke product  $B$  is uniquely determined by  $\mathcal{C}$  up to post-composition with an element of  $\text{Aut}(\mathbb{D})$ .

See [28], while the case of equality has been settled in [32]. The Blaschke product  $B$  in Theorem 3.1 is called *maximal Blaschke product for  $\mathcal{C}$* . The set of all maximal Blaschke products will be denoted by MBP.

**Remarks 3.2 (Properties of maximal Blaschke products).**

- (a) Maximal Blaschke products are indestructible:  $f \in \text{MBP}, T \in \text{Aut}(\mathbb{D}) \implies T \circ f \in \text{MBP}$ .
- (b)  $(\text{FBP} \subseteq \text{MBP})$

Maximal Blaschke products for *finite* sets  $\mathcal{C}$  are *finite* Blaschke products and vice versa, see [32, Remark 1.2 (b)]. In particular, Theorem 3.1 generalizes Theorem 1.1 and Theorem 1.2.

- (c) Any maximal Blaschke product is uniquely determined by its critical set up to postcomposition with an element of  $\text{Aut}(\mathbb{D})$ . This does not hold for general infinite Blaschke products. Neat examples are the nontrivial Frostman shifts

$$\pi_a(z) := \frac{a - \pi_0(z)}{1 - \bar{a}\pi_0(z)}, \quad a \in \mathbb{D} \setminus \{0\},$$

of the standard singular inner function

$$\pi_0(z) := \exp\left(-\frac{1+z}{1-z}\right)$$

which are Blaschke products without critical points.

- (d) The accumulation points of the critical set of a maximal Blaschke product  $B$  are exactly the accumulation points of its zero set, and  $B$  has an analytic continuation across any other point of the unit circle, see [32, Theorem 1.4 and Corollary 1.5].
- (e) The set of maximal Blaschke products is closed with respect to composition, see [32, Theorem 1.7].

**4. Zeros sets and invariant subspaces for Bergman spaces**

**Remark 4.1 (MBPs and zero sets in Bergman spaces).** Theorem 3.1 shows in particular that a set  $\mathcal{C} \subseteq \mathbb{D}$  is the critical set of a function in  $\mathcal{B}$  if and only if it is the critical set of some maximal Blaschke product. It has been shown in [28] that this is the case if and only if  $\mathcal{C}$  is the zero set of a function in the Bergman space ([14, 23])

$$A_1^2 = \left\{ \varphi : \mathbb{D} \rightarrow \mathbb{C} \text{ holomorphic} : \iint_{\mathbb{D}} (1 - |z|^2) |\varphi(z)|^2 dx dy < \infty \right\}.$$

Hence

$$\text{MBP}/\text{Aut}(\mathbb{D}) = \{\text{zero sets of } A_1^2\}. \tag{4.1}$$

This can be seen as an analogue of the classical fact that up to a rotation (= multiplication by a number  $\eta \in \mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$ ) the zero sets of functions in the Hardy space  $H^2$  are exactly the zero sets of Blaschke products,

$$\text{BP}/\mathbb{S}^1 = \{\text{zero sets of } H^2\}.$$

**Remark 4.2 (Critical sets in  $\mathcal{B}$  and singly generated invariant subspaces in Bergman spaces).** Remark 4.1 has a simple operator theoretic interpretation, cf. [14, 23] for background. A closed subspace of  $A_1^2$  is called *zero-based* if it is defined as the set of all  $A_1^2$ -functions that vanish at a prescribed set of points in  $\mathbb{D}$ . Each such subspace is *invariant*, that is, invariant w.r.t. to multiplication by  $z$ . Hence (4.1) can be trivially rewritten as

$$\text{MBP}/\text{Aut}(\mathbb{D}) = \{ \text{zero-based invariant subspaces of } A_1^2 \} .$$

In particular, if we denote by  $[H]$  the *subspace generated by a function*  $H \in A_1^2$ , that is, the minimal closed invariant subspace of  $A_1^2$  which contains  $H$ , then each zero-based subspace of  $A_1^2$  has the form  $[B']$ , meaning that it is singly generated by the derivative  $B' \in A_1^2$  of some maximal Blaschke product  $B$ . Combining this observation with the beautiful concept of asymptotic spectral synthesis of Nikol'skii [38] and a deep result of Shimorin [43] about approximation of singly-generated invariant subspaces of Bergman spaces by zero-based subspaces, O. Ivrii [27] has recently been led to the following striking conjecture

$$\text{Inner functions}/\text{Aut}(\mathbb{D}) = \{ \text{singly generated invariant subspaces of } A_1^2 \} ,$$

or in more explicit terms:

**Conjecture 4.3 (Ivrii [27]).** Any singly generated subspace of  $A_1^2$  can be generated by the derivative of an inner function. This inner function is uniquely determined up to postcomposition with a unit disk automorphism.

This conjecture can be seen as an analogue of the celebrated result of Beurling that the invariant subspaces of  $H^2$  are generated by inner functions:

$$\text{Inner functions}/\mathbb{S}^1 = \{ \text{invariant subspaces of } H^2 \} .$$

We refer to the original papers [26, 27] for details and a number of substantial results in support of Conjecture 4.3.

## 5. The strong form of the extended Nehari–Schwarz lemma at the boundary

We now return to Theorem 3.1 and discuss the case of equality at the boundary. For this purpose it is convenient to denote by

$$f^h(z) := (1 - |z|^2) \frac{|f'(z)|}{1 - |f(z)|^2}$$

the *hyperbolic derivative* of a holomorphic function  $f : \mathbb{D} \rightarrow \mathbb{D}$ , see [4, Definition 5.1]. If  $f \in \mathcal{B}$  and  $B$  is a maximal Blaschke product with  $\mathcal{C}_B \subseteq \mathcal{C}_f$ , then

$$\frac{f^h}{B^h} : \mathbb{D} \setminus \mathcal{C}_B \rightarrow \mathbb{R}$$

has a continuous extension to  $\mathbb{D}$  which will still be denoted by  $f^h/B^h$ . Theorem 3.1 (see also [32, Theorem 2.2 (b)]) implies:

(i) (Extended Nehari–Schwarz inequality)

$$\frac{f^h(z)}{B^h(z)} \leq 1 \quad \text{for all } z \in \mathbb{D};$$

(ii) (Strong form of the extended Nehari–Schwarz lemma at an interior point)

$$\frac{f^h(z)}{B^h(z)} = 1 \quad \text{for some point } z \in \mathbb{D} \iff f = T \circ B \quad \text{for some } T \in \text{Aut}(\mathbb{D}).$$

Recently, a boundary version of this interior rigidity result for functions in  $\mathcal{B}$  has been obtained in [9]:

**Theorem 5.1 (The strong form of the generalized Nehari–Schwarz lemma at the boundary).** *Let  $\mathcal{C}$  be the critical set of a non-constant function in  $\mathcal{B}$ ,  $B$  a maximal Blaschke product with critical set  $\mathcal{C}_B = \mathcal{C}$ , and  $f \in \mathcal{B}$  such that  $\mathcal{C}_f \supseteq \mathcal{C}$ . If*

$$\frac{f^h(z_n)}{B^h(z_n)} = 1 + o((1 - |z_n|)^2)$$

for some sequence  $(z_n)$  in  $\mathbb{D}$  such that  $|z_n| \rightarrow 1$ , then  $f = T \circ B$  for some  $T \in \text{Aut}(\mathbb{D})$  and  $f$  is a maximal Blaschke product.

The proof of Theorem 5.1 in [9] is based on PDE methods, in particular a Harnack-type inequality for solutions of the Gauss curvature equation, see [9] for details. This approach also yields a version of the strong form of the Ahlfors–Schwarz lemma [1, 12, 24, 36, 40, 50] at the boundary, see [9, Theorem 2.6]. The special case  $\mathcal{C} = \emptyset$  of Theorem 5.1 is the following boundary version of the strong form of the classical Schwarz–Pick lemma:

**Theorem 5.2 (The strong form of the Schwarz–Pick lemma at the boundary).** *Let  $f : \mathbb{D} \rightarrow \mathbb{D}$  be holomorphic. If*

$$f^h(z_n) = 1 + o((1 - |z_n|)^2)$$

for some sequence  $(z_n)$  in  $\mathbb{D}$  such that  $|z_n| \rightarrow 1$ , then  $f \in \text{Aut}(\mathbb{D})$ .

The error term is sharp. For  $f(z) = z^2$  we have

$$f^h(z) = \frac{2|z|}{1 + |z|^2} = 1 - \frac{(1 - |z|)^2}{1 + |z|^2} = 1 - \frac{1}{2}(1 - |z|)^2 + o((1 - |z|)^2) \quad (|z| \rightarrow 1).$$

Hence one cannot replace “little  $o$ ” by “big  $O$ ” in Theorem 5.2. Theorem 5.2 can also be deduced from the inequality

$$f^h(z) \leq \frac{f^h(0) + \frac{2|z|}{1 + |z|^2}}{1 + f^h(0) \frac{2|z|}{1 + |z|^2}} \quad \text{for all } |z| < 1, \tag{5.1}$$

which has been proved by Golusin (see [20, Theorem 3] or [21, p. 335], and independently by Yamashita [51, 52], Beardon [3], and by Beardon & Minda [4, 5] as part of their elegant work on multi-point Schwarz–Pick lemmas. With hindsight, inequality (5.1) is exactly the case  $w = 0$  in Corollary 3.7 of [4].

**Remark 5.3 (The boundary Schwarz–Pick lemma and the boundary Schwarz lemma of Burns and Krantz).** From Theorem 5.2 one can easily deduce the well-known boundary Schwarz lemma of Burns and Krantz [10], which asserts that if  $f$  is a holomorphic selfmap of  $\mathbb{D}$  such that

$$f(z) = z + o(|1 - z|^3) \quad \text{as } z \rightarrow 1, \quad (5.2)$$

then  $f(z) \equiv z$ . We refer to [9, Remark 2.2] for details.

**Remark 5.4.** Baracco, Zaitsev and Zampieri [2] have improved the boundary Schwarz lemma of Burns and Krantz by proving that if  $f : \mathbb{D} \rightarrow \mathbb{D}$  is a holomorphic map such that

$$f(z_n) = z_n + o(|1 - z_n|^3)$$

for some sequence  $(z_n)$  in  $\mathbb{D}$  converging nontangentially to 1, then  $f(z) \equiv z$ . Does the result of Baracco, Zaitsev and Zampieri follow from Theorem 5.2?

We refer to [7, 11, 13, 39, 44, 46] and in particular to the survey [17] by Elin et al. for more on boundary Schwarz–type lemmas in the one variable setting.

## References

- [1] Ahlfors, L.V., *An extension of Schwarz’s lemma*, Trans. Am. Math. Soc., **43**(1938), 359-364.
- [2] Baracco, L., Zaitsev, D., Zampieri, G., *A Burns-Krantz type theorem for domains with corners*, Math. Ann., **336**(2006), no. 3, 491-504.
- [3] Beardon, A.F., *The Schwarz-Pick Lemma for derivatives*, Proc. Am. Math. Soc., **125**(1997), no. 11, 3255-3256.
- [4] Beardon, A.F., Minda, D., *A multi-point Schwarz-Pick lemma*, J. Anal. Math., **92**(2004), 81-104.
- [5] Beardon, A.F., Minda, D., *The hyperbolic metric and geometric function theory*, In: Proceedings of the International Workshop on Quasiconformal Mappings and Their Applications, December 27, 2005-January 1, 2006, 9-56, New Delhi, Narosa Publishing House, 2007.
- [6] Bergweiler, W., *A new proof of the Ahlfors five islands theorem*, J. Anal. Math., **76**(1998), 337-347.
- [7] Bolotnikov, V., *A uniqueness result on boundary interpolation*, Proc. Am. Math. Soc., **136**(2008), no. 5, 1705-1715.
- [8] Bousch, T., *Sur quelques problèmes de dynamique holomorphe*, Ph.D. thesis, Université Paris 11, Orsay, 1992.
- [9] Bracci, F., Kraus, D., Roth, O., *A new Schwarz-Pick lemma at the boundary and rigidity of holomorphic maps*, arXiv:2003.02019.
- [10] Burns, D.M., Krantz, S., *Rigidity of holomorphic mappings and a new Schwarz lemma at the boundary*, J. Am. Math. Soc., **7**(1994), no. 3, 661-676.
- [11] Chelst, D., *A generalized Schwarz lemma at the boundary*, Proc. Am. Math. Soc., **129**(2001), no. 11, 3275-3278.
- [12] Chen, H., *On the Bloch Constant*, In: Arakelian, N. (ed.) et al., Approximation, Complex Analysis, and Potential Theory, Kluwer Academic Publishers, 2001, 129-161.



- [13] Dubinin, V.N., *The Schwarz inequality on the boundary for functions regular in the disk*, J. Math. Sci., **122**(2004), no. 6, 3623-3629.
- [14] Duren, P., Schuster, A., *Bergman Spaces*, Amer. Math. Soc., 2004.
- [15] Dyakonov, K.M., *A characterization of Möbius transformations*, C.R. Math. Acad. Sci. Paris, **352**(2014), no. 2, 593-595.
- [16] Dyakonov, K.M., *Inner functions and inner factors of their derivatives*, Integr. Equ. Oper. Theory, **82**(2015), no. 2, 151-155.
- [17] Elin, M., Jacobzon, F., Levenshtein, M., Shoikhet, D., *The Schwarz lemma: rigidity and dynamics*, In: A. Vasil'ev, Harmonic and Complex Analysis and its Applications, Springer, 2014, 135-230.
- [18] Frostman, O., *Sur les produits de Blaschke*, Fysiogr. Sällsk. Lund Förh, **12**(1942), no. 15, 169-182.
- [19] Garcia, S.R., Mashreghi, J., Ross, W.T., *Finite Blaschke Products and Their Connections*, Springer, 2018.
- [20] Golusin, G.M., *Some estimations of derivatives of bounded functions*, (Russian. English Summary), Rec. Math. [Mat. Sbornik] N.S., **16(58)**(1945), 295-306.
- [21] Golusin, G.M., *Geometric Theory of Functions of One Complex Variable*, AMS, 1969.
- [22] Graham, I., Minda, D., *A Schwarz lemma for multivalued functions and distortion theorems for Bloch functions with branch points*, Trans. Amer. Math. Soc., **351**(1999), no. 12, 4741-4752.
- [23] Hedenmalm, H., Korenblum, B., Zhu, K., *Theory of Bergman Spaces*, Springer, 2000.
- [24] Heins, M., *On a class of conformal metrics*, Nagoya Math. J., **21**(1962), 1-60.
- [25] Heins, M., *Some characterizations of finite Blaschke products of positive degree*, J. Anal. Math., **46**(1986), 162-166.
- [26] Ivrii, O., *Prescribing inner parts of derivatives of inner functions*, J. Anal. Math., **139**(2019), no. 2, 495-519.
- [27] Ivrii, O., *Critical structures of inner functions*, J. Funct. Anal., **281**(2021), no. 8, Paper No. 109138, 14 pp.
- [28] Kraus, D., *Critical sets of bounded analytic functions, zero sets of Bergman spaces and nonpositive curvature*, Proc. London Math. Soc., **106**(2013), no. 3, 931-956.
- [29] Kraus, D., Roth, O., *Critical points of inner functions, nonlinear partial differential equations, and an extension of Liouville's theorem*, J. Lond. Math. Soc., **77**(2008), no. 1, 183-202.
- [30] Kraus, D., Roth, O., *The behaviour of solutions of the Gaussian curvature equation near an isolated boundary point*, Math. Proc. Cambridge Philos. Soc., **145**(2008), no. 3, 643-667.
- [31] Kraus, D., Roth, O., *Critical points, the Gauss curvature equation and Blaschke products*, Fields Institute Comm., **65**(2012), 133-157.
- [32] Kraus, D., Roth, O., *Maximal Blaschke products*, Adv. Math., **241**(2013), 58-78.
- [33] Kraus, D., Roth, O., Ruscheweyh, St., *A boundary version of Ahlfors' Lemma, locally complete conformal metrics and conformally invariant reflection principles for analytic maps*, J. Anal. Math., **101**(2007), 219-256.
- [34] Liu, X., Minda, D., *Distortion theorems for Bloch functions*, Trans. Amer. Math. Soc., **333**(1992), no. 1, 325-338.

- [35] Minda, D., *Lower bounds for the hyperbolic metric in convex regions*, Rocky Mountain J. Math., **13**(1983), no. 1, 61-69.
- [36] Minda, D., *The strong form of Ahlfors' lemma*, Rocky Mt. J. Math., **17**(1987), 457-461.
- [37] Nehari, Z., *A generalization of Schwarz' lemma*, Duke Math. J., **14**(1947), 1035-1049.
- [38] Nikol'skii, N.K., *Treatise on the Shift Operator*, Springer 1986.
- [39] Osserman, R., *A sharp Schwarz inequality on the boundary*, Proc. Am. Math. Soc., **128**(2000), no. 12, 3513-3517.
- [40] Royden, H., *The Ahlfors-Schwarz lemma: The case of equality*, J. Anal. Math., **46**(1986), 261-270.
- [41] Semmler, G., Wegert, E., *Finite Blaschke products with prescribed critical points, Stieltjes polynomials, and moment problems*, Anal. Math. Phys., **9**(2019), no. 1, 221-249.
- [42] Shapiro, J.H., *Composition Operators and Classical Function Theory*, New York, Springer-Verlag, 1993.
- [43] Shimorin, S.M., *Approximate spectral synthesis in the Bergman space*, Duke Math. J., **101**(2000), no. 1, 1-39.
- [44] Shoikhet, D., *Another look at the Burns-Krantz theorem*, J. Anal. Math., **105**(2000), 19-42.
- [45] Stephenson, K., *Introduction to Circle Packing: The Theory of Discrete Analytic Functions*, Cambridge Univ. Press, 2005.
- [46] Tauraso, R., Vlacci, F., *Rigidity at the boundary for holomorphic self-maps of the unit disk*, Complex Variables, Theory Appl., **45**(2001), no. 2, 151-165.
- [47] Walsh, J., *The Location of Critical Points of Analytic and Harmonic Functions*, Am. Math. Soc., 1950.
- [48] Wang, Q., Peng, J., *On critical points of finite Blaschke products and the equation  $\Delta u = e^{2u}$* , (Chinese), Kexue Tongbao, **24**(1979), 583-586.
- [49] Wegert, E., *Seeing the monodromy group of a Blaschke product*, Notices Amer. Math. Soc., **67**(2020), no. 7, 965-975.
- [50] Yamada, A., *Bounded analytic functions and metrics of constant curvature on Riemann surfaces*, Kodai Math. J., **11**(1988), no. 3, 317-324.
- [51] Yamashita, S., *The Pick version of the Schwarz lemma and comparison of the Poincaré densities*, Ann. Acad. Sci. Fenn., Ser. A I, Math., **19**(1994), no. 2, 291-322.
- [52] Yamashita, S., *Goluzin's extension of the Schwarz-Pick inequality*, J. Inequal. Appl., **1**(1997), no. 4, 345-356.
- [53] Zakeri, S., *On critical points of proper holomorphic maps on the unit disk*, Bull. London Math. Soc., **30**(1996), no. 1, 62-66.

Oliver Roth

Department of Mathematics, University of Würzburg,  
Emil Fischer Strasse 40, 97074, Würzburg, Germany  
e-mail: roth@mathematik.uni-wuerzburg.de